

# ENRODEM 10%

## ORAL SOLUTION FOR POULTRY



**NAME OF THE VETERINARY MEDICINAL PRODUCT: ENRODEM 10%**, as enrofloxacin 100 mg/ml, oral solution for poultry.

**STATEMENT OF THE ACTIVE SUBSTANCE AND OTHER INGREDIENTS**

1 ml solution contains: Enrofloxacin 100 mg., Excipients (potassium hydroxide, hypromellose, benzyl alcohol, purified water) ad...1 ml

**INDICATIONS**

**In poultry:** for the treatment of gastrointestinal and respiratory infections produced by microorganisms susceptible to enrofloxacin: Gram Negative bacteria (*Pseudomonas aeruginosa.*, *Pasteurella* spp, *Salmonella* spp, *E.coli*, *Actinobacillus* spp, *Campylobacter* etc) Gram positive bacteria (some strains of *Streptococcus* and *Staphylococcus* spp.), *Mycoplasma*, *Chlamydia*.

**TARGET SPECIES:** Poultry.

**DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION**

**In poultry:** the product is administered via drinking water in dosage of 10 mg enrofloxacin/kg b.w. corresponding to 0.1 ml Enrodem 10%/kg body weight / day for 5 consecutive days. In infections with *Pseudomonas aeruginosa* the dosage is 12 mg enrofloxacin/kg body weight /day (0.12 ml Enrodem 10%/kg body weight) for 5 consecutive days. Based on the recommended doses and the number and weight of the animals to be treated, the exact daily amount of Enrodem 10% should be calculated according to the following formula:

$$\frac{\text{ml Enrodem 10\% / kg body weight/day} \times \text{mean body weight of animals to be treated (kg)}}{\text{Mean daily water consumption (liters) / animal}} = \frac{\text{ml Enrodem 10\%}}{\text{liter drinking water}}$$

The uptake of medicated water is dependant on the clinical condition of animals. In order to obtain the correct dosage, the concentration in drinking water may have to be adjusted.

**ADVICE ON CORRECT ADMINISTRATION:** During treatment, the animals should drink only medicated water. Fresh medicated water should be prepared every 24 hours. To ensure a correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

**WITHDRAWAL PERIOD:** Edible tissues of poultry: 7 days after cessation of oral medication. Not permitted for use in laying bird producing eggs for human consumption.

**SPECIAL PRECAUTIONS FOR STORAGE:** Keep away from reach and sight of children! The product should be stored in original packaging at temperatures bellow 25°C, protected from direct sunlight and freeze. Shelf life of the veterinary medicinal product as packaged for sale: 2 years. Shelf life after dilution in drinking water, according to directions: 24 hours. Shelf-life after first opening of package: 28 days. Close tight the recipient after every use. Do not use after expiry date which is stated on the label.

**SPECIAL PRECAUTIONS FOR THE DISPOSAL OF UNUSED PRODUCT OR WASTE MATERIALS**

For the environmental protection, medicines should not be disposed of via wastewater or household waste. Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

**DATE ON WHICH THE PACKAGE LEAFLET WAS LAST APPROVED:**

**OTHER INFORMATION:** PE-HD flasks of 100 ml, 500 ml, and 1000 ml. PE-HD cans of 5 l, 10 l, and 20 l. Not all pack sizes may be marketed.